

Floxuridine

About This Medicine

Floxuridine is used to treat cancer. It is given continuously in an artery (intra-arterial) through an infusion pump.

Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak, and raise your risk of bleeding.
- Nausea and vomiting (throwing up)
- Diarrhea (loose bowel movements)
- Inflammation (swelling) of your small intestine
- Soreness of the mouth and throat. You may have red areas, white patches, or sores that hurt.
- Changes in your liver function, which can cause liver failure
- Skin redness and rash
- Hair loss is often temporary, although with certain medicine, hair loss can sometimes be permanent. Hair loss may happen suddenly or gradually. If you lose hair, you may lose it from your head, face, armpits, pubic area, chest, and/or legs. You may also notice your hair getting thin.

Note: All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your medical team.

Warnings and Precautions

- Severe bone marrow suppression
- Risk of heart attack
- Severe nausea, vomiting, diarrhea and soreness of the mouth and throat
- Abnormal bleeding – symptoms may be coughing up blood, throwing up blood (may look like coffee grounds), red or black tarry bowel movements, abnormally heavy menstrual flow, nosebleeds, or any other unusual bleeding.

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.

Important Information

- Cytotoxic medicines leave the body through urine and stool, but they can also be present in other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your health care provider after your treatment. If you take a cytotoxic pill each day, follow these precautions every day.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your doctor or nurse tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your nurse before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- Drink enough fluids to keep your urine pale yellow.
- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your nurse or doctor about medicine that can lessen or stop your diarrhea.
- Mouth care is very important. Your mouth care should consist of routine, gentle cleaning of your teeth or dentures and rinsing your mouth with a mixture of 1/2 teaspoon of salt in 8 ounces of water or 1/2 teaspoon of baking soda in 8 ounces of water. This should be done at least after each meal and at bedtime.
- If you have mouth sores, avoid mouthwash that has alcohol. Also avoid alcohol and smoking because they can bother your mouth and throat.
- To help with hair loss, wash with a mild shampoo and avoid washing your hair every day. Avoid coloring your hair.
- Avoid rubbing your scalp, pat your hair or scalp dry.
- Limit your use of hair spray, electric curlers, blow dryers, and curling irons.
- If you are interested in getting a wig, talk to your nurse and they can help you get in touch with programs in your local area.



- If you get a rash do not put anything on it unless your doctor or nurse says you may. Keep the area around the rash clean and dry. Ask your doctor for medicine if your rash bothers you.

Food and Medicine Interactions

- There are no known interactions of floxuridine with food.
- Check with your doctor or pharmacist about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known medicine interactions with floxuridine. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Chest pain or symptoms of a heart attack. Most heart attacks involve pain in the center of the chest that lasts more than a few minutes. The pain may go away and come back or it can be constant. It can feel like pressure, squeezing, fullness, or pain. Sometimes pain is felt in one or both arms, the back, neck, jaw, or stomach. **If any of these symptoms last 2 minutes, call 911.**
- Tiredness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Pain in your mouth or throat that makes it hard to eat or drink
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Signs of abnormal bleeding such as coughing up blood, throwing up blood (may look like coffee grounds), easy bleeding or bruising, red or black tarry bowel movements, abnormally heavy menstrual flow, nosebleeds, or any other unusual bleeding
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- A new rash and/or itching or a rash that is not relieved by prescribed medicines
- If you think you may be pregnant

Reproduction Warnings

- **Pregnancy warning:** This medicine may have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment. Let your doctor know right away if you think you may be pregnant.
- **Breastfeeding warning:** It is not known if this medicine passes into breast milk. For this reason, women should not breastfeed during treatment because this medicine could enter the breast milk and cause harm to a breastfeeding baby.



- **Fertility warning:** Fertility studies have not been done with this medicine. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

Revised November 2023

