

Chlorambucil (Leukeran)

About This Medicine

Chlorambucil is used to treat cancer. It is given orally (by mouth).

Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak, and raise your risk of bleeding.
- Soreness of the mouth and throat. You may have red areas, white patches, or sores that hurt.
- Nausea and vomiting (throwing up)
- Diarrhea (loose bowel movements)
- Fever
- In women, menstrual bleeding may become irregular or stop while you are getting this medicine. Do not assume that you cannot become pregnant if you do not have a menstrual period.
- Effects on the nerves are called peripheral neuropathy. You may feel numbness, tingling, or pain in your hands and feet. It may be hard for you to button your clothes, open jars, or walk as usual. The effect on the nerves may get worse with more doses of the medicine. These effects get better in some people after the medicine is stopped but it does not get better in all people.
- Changes in your central nervous system can happen which can be life-threatening. The central nervous system is made up of your brain and spinal cord. You could feel extreme tiredness, agitation, confusion, have hallucinations (see or hear things that are not there), trouble understanding or speaking, loss of control of your bowels or bladder, eyesight changes, numbness or lack of strength to your arms, legs, face, or body, seizures or coma. **If you start to have any of these symptoms let your doctor know right away.**
- Rash and allergic skin reactions

Note: All possible side effects are not included. Your side effects may be different depending on your cancer diagnosis, condition, or if you are receiving other medicines in combination. Please discuss any concerns or questions with your medical team.

Warnings and Precautions

- This medicine may raise your risk of getting a second cancer, such as acute leukemia.
- Seizure. Common symptoms of a seizure can include confusion, blacking out, passing out, loss of hearing or vision, blurred vision, unusual smells or tastes (such as burning rubber), trouble talking,

tremors or shaking in parts or all of the body, repeated body movements, tense muscles that do not relax, and loss of control of urine and bowels. There are other less common symptoms of seizures.

If you or your family member suspects you are having a seizure, call 911 right away.

- Severe bone marrow suppression
- Changes in your liver function
- Severe allergic skin reaction. You may develop blisters on your skin that are filled with fluid or a severe red rash all over your body that may be painful.
- Thickening and/or inflammation (swelling) of the lung tissues. You may have a dry cough or trouble breathing.

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.

Important Information

- Talk to your doctor before receiving any vaccinations during your treatment. Some vaccinations are not recommended while receiving chlorambucil.

How to Take Your Medication

- Swallow the medicine whole without food.
- **Missed dose:** If you vomit or miss a dose, contact your doctor for further instructions. Do not take 2 doses at the same time or extra doses.
- **Handling:** Wash your hands with soap and water, before and after handling your medicine, your caretakers should not handle your medicine with bare hands and should wear latex gloves.
- Cytotoxic medicines leave the body through urine and stool, but they can also be present in other body fluids such as blood, vomit, semen, and vaginal fluids. Take precautions to prevent others from coming in contact with your medicine or your body fluids. Follow safety precautions during your treatment and for as long as directed by your health care provider after your treatment. If you take a cytotoxic pill each day, follow these precautions every day.
- **Storage:** Store this medicine in the refrigerator, between 36°F to 46°F (2°C to 8°C).
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medication label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your nurse or your doctor how to dispose of unused medicine.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your doctor or nurse tells you, and whenever you feel like you may have a fever.



- To help decrease the risk of bleeding, use a soft toothbrush. Check with your nurse before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Drink enough fluids to keep your urine pale yellow.
- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your nurse or doctor about medicine that can lessen or stop your diarrhea.
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- Mouth care is very important. Your mouth care should consist of routine, gentle cleaning of your teeth or dentures and rinsing your mouth with a mixture of 1/2 teaspoon of salt in 8 ounces of water or 1/2 teaspoon of baking soda in 8 ounces of water. This should be done at least after each meal and at bedtime.
- If you have mouth sores, avoid mouthwash that has alcohol. Also avoid alcohol and smoking because they can bother your mouth and throat.
- If you get a rash do not put anything on it unless your doctor or nurse says you may. Keep the area around the rash clean and dry. Ask your doctor for medicine if your rash bothers you.
- If you have numbness and tingling in your hands and feet, be careful when cooking, walking, and handling sharp objects and hot liquids.

Food and Medicine Interactions

- There are no known interactions of chlorambucil with food, however this medication should be taken on an empty stomach.
- This medicine may interact with other medicines. Tell your doctor and pharmacist about all the prescription and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) that you are taking at this time. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Easy bleeding or bruising
- Extreme tiredness, agitation, or confusion



- Hallucinations
- Trouble understanding or speaking
- Blurry vision or changes in your eyesight
- Symptoms of a seizure such as confusion, blacking out, passing out, loss of hearing or vision, blurred vision, unusual smells or tastes (such as burning rubber), trouble talking, tremors or shaking in parts or all of the body, repeated body movements, tense muscles that do not relax, and loss of control of urine and bowels. **If you or your family member suspects you are having a seizure, call 911 right away.**
- Pain in your chest
- Dry cough
- Trouble breathing
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Pain in your mouth or throat that makes it hard to eat or drink
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- New rash and/or itching
- Rash that is not relieved by prescribed medicines
- Unusual skin lumps/bumps
- Irregular and/or no menstrual bleeding
- Flu-like symptoms: fever, headache, muscle and joint aches, and fatigue (low energy, feeling weak)
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Numbness, tingling, or pain in your hands and feet
- Loss of control of bowels or bladder
- Numbness or lack of strength to your arms, legs, face, or body
- If you think you may be pregnant

Reproduction Warnings

- **Pregnancy warning:** This medicine can have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment. Let your doctor know right away if you think you may be pregnant.
- **Breastfeeding warning:** It is not known if this medicine passes into breast milk. For this reason, women should talk to their doctor about the risks and benefits of breastfeeding during treatment with this medicine because this medicine may enter the breast milk and cause harm to a breastfeeding baby
- **Fertility warning:** In men and women both, this medicine may affect your ability to have children in the future. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

Revised November 2023

